

## 國立臺灣科技大學 110 學年度碩士班招生試題

系所組別：工業管理系碩士班甲組、乙組、丙組  
 科目：統計學

(總分為 100 分)

(Total 100 Points.) *There are 7 problems in this exam. Show intermediate steps and formulas for partial credit. You must explain how you compute your results or answers for full credit.*

- (20 points) There are 300 passengers in an international flight. We know that 150, 100, and 50 passengers in this flight are from countries A, B, and C, respectively. Each passenger may carry COVID-19 virus with a certain probability independently. These probabilities for the passengers from countries A, B, and C are 3%, 4%, and 5%, respectively.
  - If 5 passengers are selected at random from country C passengers, find the probability that there is exactly 1 passenger carrying COVID-19 virus. (5 points)
  - If a passenger is selected at random from the flight, find the probability that this passenger is carrying COVID-19 virus. (5 points)
  - If 5 passengers are selected at random from the flight, find the probability that there is exactly 1 passenger carrying COVID-19 virus. (5 points)
  - Suppose that a passenger is selected at random from the flight and is found to carry COVID-19 virus. Find the probability that the passenger is from country C. (5 points)
- (20 points) Three balls are drawn without replacement from 12 balls (4 white, 4 black, and 4 red). Let  $X$  be the number of white balls selected and  $Y$  be the number of black balls selected.
  - Find the joint probability distribution of  $X$  and  $Y$ . (5 points)
  - Find the probability  $\mathbb{P}\{Y = 1\}$ . (5 points)
  - Find the conditional distribution of  $X$ , given that  $Y = 1$ . (5 points)
  - Find the probability  $\mathbb{P}\{X + Y \geq 2\}$ . (5 points)
- (10 points) If the joint density function of two random variables  $X$  and  $Y$  is given by  $f(x, y) = 2$  for  $0 < x \leq y < 1$ . What is the correlation coefficient of  $X$  and  $Y$ ?
- (15 points) A company that is looking at the time to failure of the parts (days) it uses decides to look at an alternative supplier of the product. The data for the current supplier and for the new supplier are shown here.

	1	2	3	4	5	6	7	8	9
Current Supplier	130	100	110	100	70	100	90		
Alternative Supplier	120	100	120	140	100	140	110	130	120

Assume that the data are normally distributed.

- Please test whether the variances of the time to failure of the parts produced by different suppliers are equal. (Please clearly state the null, the alternative hypothesis, and the test statistic, and use  $\alpha = 0.10$ ) (5 points)
- The company has decided that if the mean time to failure for the new suppliers is significantly longer than it is for the current supplier, it will switch suppliers. Should the company switch suppliers? (Please clearly state the null, the alternative hypothesis, and the test statistic, and use  $\alpha = 0.05$ ) (10 points)



## 國立臺灣科技大學 110 學年度碩士班招生試題

系所組別：工業管理系碩士班甲組、乙組、丙組

科 目：統計學

(總分為 100 分)

5. (15 points) Three different machines are being considered for use in manufacturing rubber seals. The machines are being compared with respect to tensile strength of the product. A random sample of five seals from each machine is used to determine whether the mean tensile strength varies from machine to machine. The company want to know whether or not the mean tensile strength is the same for all three machines.

Machine		
A	B	C
20	17	22
22	19	20
20	19	20
18	20	21
20	20	22

- (a) Please construct the ANOVA table. (10 points)
- (b) Please state the null and alternative hypothesis, perform the test, and state your conclusion using 0.05 level of significance. (5 points)
6. (10 points) A real estate economist collects data on two similar neighborhoods, one bordering a MRT station, and one that is a neighborhood about 3 kilometers from the MRT station. She records 50 observations of real estate values, where house prices are given in ten thousand NT\$ (i.e., NT\$ 10000), and size is the number of square meter of living area.

Consider the partial printout for a regression analysis of the relationship between prices (in ten thousands) and two independent variables Size and MRT. As a house can only be near the MRT station or not, the MRT variable is coded as

$$\text{MRT} = \begin{cases} 1 & \text{near the MRT station} \\ 0 & \text{otherwise} \end{cases}$$

Predictor	Coef	SE	T	P
Constant	150	50	3	0.0043
Size	10	1	10	0.0000
MRT	60	25	2.4	0.0205
Size*MRT	5	2	2.5	0.0160

Source	SS	df	MS	F	P
Regression	72000	3	24000	15	<0.0001
Residual	73600	46	1600		
Total	145600	49			

- (a) Plot the least squares prediction equation associated with two different neighborhoods, and discuss your findings. (5 points)
- (b) Test the overall utility of the model (Please clearly state the null, the alternative hypothesis, and the test statistic, and use  $\alpha = 0.05$ ). (5 points)
7. (10 points) Let  $X_1, X_2, \dots, X_n$  denote a random sample from a distribution that is  $N(\theta, 1)$ , where the mean  $\theta$  is unknown. Please show that there is no uniformly most powerful (UMP) test of the simple hypothesis  $H_0 : \theta = \theta_0$ , where  $\theta_0$  be a specified value of  $\theta$ , against the alternative hypothesis  $H_1 : \theta \neq \theta_0$ .




國立臺灣科技大學 110 學年度碩士班招生試題

系所組別：工業管理系碩士班甲組、乙組、丙組  
 科目：統計學

(總分為 100 分)

Table A.4 Student *t*-Distribution Probability Table




<i>v</i>	$\alpha$							
	0.40	0.30	0.20	0.15	0.10	0.05	0.025	0.01
1	0.325	0.727	1.376	1.963	3.078	6.314	12.706	31.821
2	0.289	0.617	1.061	1.386	1.886	2.920	4.303	6.965
3	0.277	0.584	0.978	1.250	1.638	2.353	3.182	5.041
4	0.271	0.569	0.941	1.190	1.533	2.132	2.776	4.088
5	0.267	0.559	0.920	1.156	1.476	2.015	2.571	3.365
6	0.265	0.553	0.906	1.134	1.440	1.943	2.447	3.143
7	0.263	0.549	0.896	1.119	1.415	1.895	2.365	2.998
8	0.262	0.546	0.889	1.108	1.397	1.860	2.306	2.896
9	0.261	0.543	0.883	1.100	1.383	1.833	2.262	2.821
10	0.260	0.542	0.879	1.093	1.372	1.812	2.228	2.764
11	0.260	0.540	0.876	1.088	1.363	1.796	2.201	2.718
12	0.259	0.539	0.873	1.083	1.356	1.782	2.179	2.681
13	0.259	0.538	0.870	1.079	1.350	1.771	2.160	2.650
14	0.258	0.537	0.868	1.076	1.345	1.761	2.145	2.624
15	0.258	0.536	0.866	1.074	1.341	1.753	2.131	2.602
16	0.258	0.535	0.865	1.071	1.337	1.746	2.120	2.583
17	0.257	0.534	0.863	1.069	1.333	1.740	2.110	2.567
18	0.257	0.534	0.862	1.067	1.330	1.734	2.101	2.552
19	0.257	0.533	0.861	1.066	1.328	1.729	2.093	2.539
20	0.257	0.533	0.860	1.064	1.325	1.725	2.086	2.528
21	0.257	0.532	0.859	1.063	1.323	1.721	2.080	2.518
22	0.256	0.532	0.858	1.061	1.321	1.717	2.074	2.508
23	0.256	0.532	0.858	1.060	1.319	1.714	2.069	2.500
24	0.256	0.531	0.857	1.059	1.318	1.711	2.064	2.492
25	0.256	0.531	0.856	1.058	1.316	1.708	2.060	2.485
26	0.256	0.531	0.856	1.058	1.315	1.706	2.052	2.479
27	0.256	0.531	0.855	1.057	1.314	1.703	2.052	2.473
28	0.256	0.530	0.855	1.056	1.313	1.701	2.048	2.467
29	0.256	0.530	0.854	1.055	1.311	1.699	2.045	2.462
30	0.256	0.530	0.854	1.055	1.310	1.697	2.042	2.457
40	0.255	0.529	0.851	1.050	1.303	1.684	2.021	2.423
60	0.254	0.527	0.848	1.045	1.296	1.671	2.000	2.390
120	0.254	0.526	0.845	1.041	1.289	1.658	1.980	2.358
$\infty$	0.253	0.524	0.842	1.036	1.282	1.645	1.960	2.326

737

738

Appendix A Statistical Tables and Proofs

<i>v</i>	$\alpha$									
	0.02	0.015	0.01	0.0075	0.005	0.0025	0.001	0.0005	0.00025	0.0001
1	15.894	21.205	31.821	42.433	63.656	127.321	636.578	1273.21	31600	636578
2	4.849	5.643	6.965	8.073	9.925	14.089	31.600	63.656	127.321	316.000
3	3.482	3.896	4.541	5.047	5.841	7.453	12.924	20.130	30.829	63.656
4	2.999	3.298	3.747	4.088	4.604	5.598	8.610	12.924	20.130	30.829
5	2.757	3.003	3.365	3.634	4.032	4.773	6.869	10.591	15.086	20.130
6	2.612	2.829	3.143	3.372	3.707	4.317	5.959	8.591	12.924	20.130
7	2.517	2.715	2.998	3.203	3.499	4.029	5.408	7.453	10.591	15.086
8	2.449	2.634	2.896	3.085	3.355	3.833	5.041	6.869	9.925	14.089
9	2.398	2.574	2.821	2.998	3.250	3.690	4.781	5.841	8.073	9.925
10	2.359	2.527	2.764	2.932	3.169	3.581	4.587	5.408	7.453	9.925
11	2.328	2.491	2.718	2.879	3.106	3.497	4.437	5.041	6.869	9.925
12	2.303	2.461	2.681	2.836	3.055	3.428	4.318	4.841	6.314	8.073
13	2.282	2.436	2.650	2.801	3.012	3.372	4.221	4.641	5.841	7.453
14	2.264	2.415	2.624	2.771	2.977	3.326	4.140	4.473	5.408	6.869
15	2.249	2.397	2.602	2.746	2.947	3.286	4.073	4.318	5.041	6.314
16	2.235	2.382	2.583	2.724	2.921	3.252	4.015	4.174	4.841	5.841
17	2.224	2.368	2.567	2.706	2.898	3.222	3.965	4.032	4.641	5.408
18	2.214	2.356	2.552	2.689	2.878	3.197	3.922	3.922	4.473	5.041
19	2.205	2.346	2.539	2.674	2.861	3.174	3.883	3.883	4.318	4.841
20	2.197	2.336	2.528	2.661	2.845	3.153	3.850	3.850	4.174	4.641
21	2.189	2.328	2.518	2.649	2.831	3.135	3.819	3.819	4.032	4.473
22	2.183	2.320	2.508	2.639	2.819	3.119	3.792	3.792	3.922	4.318
23	2.177	2.313	2.500	2.629	2.807	3.104	3.768	3.768	3.819	4.174
24	2.172	2.307	2.492	2.620	2.797	3.091	3.745	3.745	3.725	4.032
25	2.167	2.301	2.485	2.612	2.787	3.078	3.725	3.725	3.634	3.883
26	2.162	2.296	2.479	2.605	2.779	3.067	3.707	3.707	3.581	3.819
27	2.158	2.291	2.473	2.598	2.771	3.057	3.689	3.689	3.541	3.768
28	2.154	2.286	2.467	2.592	2.763	3.047	3.674	3.674	3.501	3.725
29	2.150	2.282	2.462	2.586	2.756	3.038	3.660	3.660	3.460	3.689
30	2.147	2.278	2.457	2.581	2.750	3.030	3.646	3.646	3.421	3.650
40	2.123	2.250	2.423	2.542	2.704	2.971	3.551	3.551	3.373	3.600
60	2.099	2.223	2.390	2.504	2.660	2.915	3.460	3.460	3.326	3.551
120	2.076	2.196	2.358	2.468	2.617	2.860	3.373	3.373	3.286	3.501
$\infty$	2.054	2.170	2.326	2.432	2.576	2.807	3.290	3.290	3.246	3.451



國立臺灣科技大學 110 學年度碩士班招生試題


系所組別：工業管理系碩士班甲組、乙組、丙組  
 科目：統計學

(總分為 100 分)

741

Table A.6 Critical Values of the F-Distribution

$f_{\alpha}(v_1, v_2)$



$v_2$	$v_1$								
	1	2	3	4	5	6	7	8	9
1	161.45	199.50	215.71	224.58	230.16	233.99	236.77	238.88	240.54
2	18.51	19.00	19.16	19.25	19.30	19.33	19.35	19.37	19.38
3	10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81
4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00
5	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77
6	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10
7	5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68
8	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39
9	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18
10	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02
11	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90
12	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80
13	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71
14	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65
15	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59
16	4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54
17	4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49
18	4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46
19	4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42
20	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39
21	4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.37
22	4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34
23	4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32
24	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30
25	4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28
26	4.23	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27
27	4.21	3.35	2.96	2.73	2.57	2.46	2.37	2.31	2.25
28	4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24
29	4.18	3.33	2.93	2.70	2.55	2.43	2.35	2.28	2.22
30	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21
40	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12
60	4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04
120	3.92	3.07	2.68	2.45	2.29	2.18	2.09	2.02	1.96
$\infty$	3.84	3.00	2.60	2.37	2.21	2.10	2.01	1.94	1.88

Reproduced from Table 18 of *Biometrika Tables for Statisticians*, Vol. 1, by permission of E.S. Pearson and the Biometrika Trustees.

742

Table A.6 (continued) Critical Values of the F-Distribution

$f_{\alpha}(v_1, v_2)$

Appendix A Statistical Tables and Proofs

$v_2$	$v_1$														
	10	12	15	20	24	30	40	60	120	$\infty$					
1	241.88	243.91	245.95	248.01	249.05	250.10	251.14	252.20	253.25	254.31					
2	19.40	19.41	19.43	19.45	19.45	19.46	19.47	19.48	19.49	19.50					
3	8.79	8.74	8.70	8.66	8.64	8.62	8.59	8.57	8.55	8.53					
4	5.96	5.91	5.86	5.80	5.77	5.75	5.72	5.69	5.66	5.63					
5	4.74	4.68	4.62	4.56	4.53	4.50	4.46	4.43	4.40	4.36					
6	4.06	4.00	3.94	3.87	3.84	3.81	3.77	3.74	3.70	3.67					
7	3.64	3.57	3.51	3.44	3.41	3.38	3.34	3.30	3.27	3.23					
8	3.35	3.28	3.22	3.15	3.12	3.08	3.04	3.01	2.97	2.93					
9	3.14	3.07	3.01	2.94	2.90	2.86	2.83	2.79	2.75	2.71					
10	2.98	2.91	2.85	2.77	2.74	2.70	2.66	2.62	2.58	2.54					
11	2.85	2.79	2.72	2.65	2.61	2.57	2.53	2.49	2.45	2.40					
12	2.75	2.69	2.62	2.54	2.51	2.47	2.43	2.38	2.34	2.30					
13	2.67	2.60	2.53	2.46	2.42	2.38	2.34	2.29	2.25	2.21					
14	2.60	2.53	2.46	2.39	2.35	2.31	2.27	2.22	2.18	2.13					
15	2.54	2.48	2.40	2.33	2.29	2.25	2.20	2.16	2.11	2.07					
16	2.49	2.42	2.35	2.28	2.24	2.19	2.15	2.11	2.06	2.01					
17	2.45	2.38	2.31	2.23	2.19	2.15	2.10	2.06	2.01	1.96					
18	2.41	2.34	2.27	2.19	2.15	2.11	2.06	2.02	1.97	1.92					
19	2.38	2.31	2.23	2.16	2.11	2.07	2.03	1.98	1.93	1.88					
20	2.35	2.28	2.20	2.12	2.08	2.04	1.99	1.95	1.90	1.84					
21	2.32	2.25	2.18	2.10	2.05	2.01	1.96	1.92	1.87	1.81					
22	2.30	2.23	2.15	2.07	2.03	1.98	1.94	1.89	1.84	1.78					
23	2.27	2.20	2.13	2.05	2.01	1.96	1.91	1.86	1.81	1.76					
24	2.25	2.18	2.11	2.03	1.98	1.94	1.89	1.84	1.79	1.73					
25	2.24	2.16	2.09	2.01	1.96	1.92	1.87	1.82	1.77	1.71					
26	2.22	2.15	2.07	1.99	1.95	1.90	1.85	1.80	1.75	1.69					
27	2.20	2.13	2.06	1.97	1.93	1.88	1.84	1.79	1.73	1.67					
28	2.19	2.12	2.04	1.96	1.91	1.87	1.82	1.77	1.71	1.65					
29	2.18	2.10	2.03	1.94	1.90	1.85	1.81	1.75	1.70	1.64					
30	2.16	2.09	2.01	1.93	1.89	1.84	1.79	1.74	1.68	1.62					
40	2.08	2.00	1.92	1.84	1.79	1.74	1.69	1.64	1.58	1.51					
60	1.99	1.92	1.84	1.75	1.70	1.65	1.59	1.53	1.47	1.39					
120	1.91	1.83	1.75	1.66	1.61	1.55	1.50	1.43	1.35	1.25					
$\infty$	1.83	1.75	1.67	1.57	1.52	1.46	1.39	1.32	1.22	1.00					

