

國立台灣科技大學九十五學年度碩士班招生試題

系所組別：財務金融研究所碩士班

科目：統計學

「總分 100 分」

25分 1. Consider the function of two variables

$$f_{XY}(x, y) = \begin{cases} Axy, & 0 < x < y, \quad 0 < y < 1 \\ 0, & \text{otherwise} \end{cases}$$

- (a) Find A such that this is a proper p.d.f.
- (b) Find the probability that $0 < x < 0.4$ and $0.6 < y < 1$
- (c) Obtain the marginal p.d.f.'s for X and Y .

25分 2. Many companies must accurately estimate their costs before a job is begun in order to acquire a contract and make a profit. For example, a heating and plumbing contractor may estimate costs for new homes based on the total area of the house and whether central air conditioning is to be installed.

- (a) Write a main effects model relating the mean cost of material and labor, $E(y)$, to the area and central-air-conditioning variables.
- (b) Write a complete second-order model for the mean cost as a function of the same two variables.
- (c) The contractor samples 25 recent jobs, fitting both the complete second-order model (part b) and the reduced main effects model (part a), so that a test can be conducted to determine whether the additional complexity of the second-order model is necessary. The resulting SSE and R^2 values are shown in the table below. Is there sufficient evidence to conclude that the second-order terms are important for predicting the mean cost? Use $\alpha = 0.05$. (There is no probability table available, please write in more detail.)

	SSE	R^2
Main effects	8.548	0.950
Second-order	6.133	0.964

173



國立台灣科技大學九十五學年度碩士班招生試題

系所組別：財務金融研究所碩士班

科目：統計學

3. Samantha was a student at University of Wisconsin - Stout majoring in Accounting. In the summer after her sophomore year she got an intern job at the First National Bank. As her boss, Brad, assumed that she must have taken statistics course in the business school, he asked Samantha to give him a 95% interval estimate of the average amount of money their customers opened a certificate deposit with the bank. Samantha randomly selected 5 from all the certificate deposits currently valid with them and found the principals were \$5,800, \$6,000, \$5,300, \$8,100, and 7,300. Samantha gave an interval estimate of between \$5,292 to \$7,708. Was she correct? If not, what do you think a correct interval would be? (16 points)
4. The manager of an auditorium where pop concerts are often held has found that the attendance to this kind of concert is normally distributed with an average of 6,000 people, and standard deviation 1,500. The tickets are sold at a flat rate of \$500 each. Overall, all concert organizers who rent this auditorium lose money on 20% of time. For the upcoming concert, how many people the concert must draw in order for the concert organizer to break even? (17 points)
5. Suppose the three most favorite kinds of food ordered at local McDonald's stores were Big Mac, French Fries, and Chicken McNuggets. To focus on those 3 kinds of food, all the customers who ordered any single selections or combinations of the three selections were called the MFM segment. A staff with an MBA degree was called to do an analysis. She reported to the COO, "Based on our internal records, 43.6% of our MFM customers ordered Big Mac, 52% bought French Fries, and 30% ate Chicken McNuggets." Proudly she continued, "My analysis shows that 1 out of 10 MFM customers ordered Big Mac and French Fries, 15.6% ordered Chicken McNuggets and French Fries. And, surprisingly, none had Big Mac and Chicken McNuggets together." The COO then asked a question, "What is the percentage of MFM customers who ordered either French Fries or Chicken McNugget alone?" A correct answer by the staff should be _____. (17 points)

※ 所有答案必須寫在
答案卷上。




國立台灣科技大學九十五學年度碩士班招生試題

系所組別：財務金融研究所碩士班

科目：統計學

TABLE 1 Values of the Standard Normal Distribution Function

$$\Phi(z) = \int_{-\infty}^z \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2}u^2\right) du = P(Z \leq z)$$



z	0	1	2	3	4	5	6	7	8	9
-3.	.0013	.0010	.0007	.0005	.0003	.0002	.0002	.0001	.0001	.0000
-2.9	.0019	.0018	.0017	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0126	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0238	.0233
-1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0300	.0294
-1.7	.0446	.0437	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0570	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-.7	.2420	.2389	.2358	.2327	.2297	.2266	.2236	.2206	.2176	.2148
-.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-.2	.4202	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-.1	.4607	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

TABLE 1 Values of the Standard Normal Distribution Function (Continued)

z	0	1	2	3	4	5	6	7	8	9
0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
.7	.7580	.7611	.7642	.7673	.7703	.7734	.7764	.7794	.7823	.7852
.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9278	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9430	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9648	.9656	.9664	.9671	.9678	.9686	.9693	.9700	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9762	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9874	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.	.9987	.9990	.9993	.9995	.9997	.9998	.9998	.9999	.9999	1.0000

Note 1: If a normal variable X is not "standard," its values must be "standardized": $Z = (X - \mu)/\sigma$, i.e., $P(X \leq x) = \Phi((x - \mu)/\sigma)$.

Note 2: For "two-tail" probabilities, see Table 1b.

Note 3: For $z \geq 4$, $\Phi(z) = 1$ to four decimal places; for $z \leq -4$, $\Phi(z) = 0$ to four decimal places.

Note 4: Entries opposite 3 and -3 are for 3.0, 3.1, 3.2, etc., and -3.0, -3.1, etc., respectively.



國立台灣科技大學九十五學年度碩士班招生試題

系所組別：財務金融研究所碩士班

科目：統計學

566

Appendix

Table 2. Percentage Points of the Chi-squared Distribution*

$1 - \alpha$	0.005	0.010	0.025	0.050	0.100	0.250	0.500	0.750	0.900	0.950	0.975	0.990	0.995
1	0.00393	0.0157	0.04952	0.07393	0.10158	0.102	0.455	1.32	2.71	3.84	5.02	6.63	7.88
2	0.01000	0.0201	0.0506	0.103	0.211	0.575	1.39	2.77	4.61	5.99	7.38	9.21	10.6
3	0.0717	0.115	0.216	0.352	0.584	1.21	2.37	4.11	6.25	7.81	9.35	11.3	12.8
4	0.207	0.297	0.484	0.711	1.06	1.92	3.36	5.39	7.78	9.49	11.1	13.3	14.9
5	0.412	0.554	0.831	1.15	1.61	2.67	4.35	6.63	9.24	11.1	12.8	15.1	16.7
6	0.676	0.872	1.24	1.64	2.20	3.45	5.35	7.84	10.6	12.6	14.4	16.8	18.5
7	0.989	1.24	1.69	2.17	2.83	4.25	6.35	9.04	12.0	14.1	16.0	18.5	20.3
8	1.34	1.65	2.18	2.73	3.49	5.07	7.34	10.2	13.4	15.5	17.5	20.1	22.0
9	1.73	2.09	2.70	3.33	4.17	5.90	8.34	11.4	14.7	16.9	19.0	21.7	23.6
10	2.16	2.66	3.25	3.94	4.87	6.75	9.34	12.5	16.0	18.3	20.5	23.2	25.2
11	2.60	3.05	3.82	4.57	5.58	7.58	10.3	13.7	17.1	19.7	21.9	24.7	26.8
12	3.07	3.57	4.40	5.23	6.30	8.44	11.3	14.8	18.5	21.0	23.3	26.2	28.3
13	3.57	4.11	5.01	5.89	7.04	9.30	12.3	16.0	19.8	22.4	24.7	27.7	29.8
14	4.07	4.66	5.63	6.57	7.79	10.2	13.3	17.1	21.1	23.7	26.1	29.1	31.3
15	4.60	5.23	6.26	7.26	8.55	11.0	14.3	18.2	22.3	25.0	27.5	30.6	32.8
16	5.14	5.81	6.91	7.96	9.31	11.9	15.3	19.4	23.5	26.3	28.8	32.0	34.3
17	5.70	6.41	7.56	8.67	10.1	12.8	16.3	20.6	24.8	27.6	30.2	33.4	35.7
18	6.28	7.01	8.23	9.39	10.9	13.7	17.3	21.9	26.0	28.9	31.5	34.8	37.2
19	6.84	7.53	8.91	10.1	11.7	14.6	18.3	22.7	27.2	30.1	32.9	36.2	38.6
20	7.43	8.26	9.59	10.9	12.4	15.5	19.3	23.8	28.4	31.4	34.2	37.6	40.0
21	8.03	8.90	10.3	11.6	13.2	16.3	20.3	24.9	29.6	32.7	35.5	38.9	41.4
22	8.64	9.54	11.0	12.3	14.0	17.2	21.3	26.0	30.8	33.9	36.8	40.3	42.8
23	9.28	10.2	11.7	13.1	14.8	18.1	22.3	27.1	32.0	35.2	38.1	41.6	44.2
24	9.89	10.9	12.4	13.8	15.7	19.0	23.3	28.2	33.2	36.4	39.4	43.0	45.6
25	10.5	11.5	13.1	14.6	16.5	19.9	24.3	29.3	34.4	37.7	40.6	44.3	46.9
26	11.2	12.2	13.8	15.4	17.3	20.8	25.3	30.4	35.6	38.9	41.9	45.6	48.3
27	11.8	12.9	14.6	16.2	18.1	21.7	26.3	31.5	36.7	40.1	43.2	47.0	49.6
28	12.5	13.6	15.3	16.9	18.9	22.7	27.3	32.6	37.9	41.3	44.5	48.3	51.0
29	13.1	14.3	16.0	17.7	19.6	23.6	28.3	33.7	39.1	42.6	45.7	49.6	52.3
30	13.8	15.0	16.8	18.5	20.6	24.5	29.3	34.8	40.3	43.8	47.0	50.9	53.7

* Abridged from Catherine M. Thompson: Tables of percentage points of the incomplete beta function and of the chi-square distribution. *Biometrika*, vol. 32 (1941), pp. 187-191, and published here with the kind permission of the editor of *Biometrika*.

Appendix 567

Table 8. Upper Percentage Points of the t Distribution*

$1 - \alpha$	0.75	0.90	0.95	0.975	0.99	0.995	0.9995
1	1.000	3.078	6.314	12.706	31.821	63.657	636.619
2	0.816	1.886	2.920	4.303	6.965	9.925	31.598
3	0.765	1.638	2.353	3.182	4.541	5.841	12.941
4	0.741	1.533	2.132	2.776	3.747	4.604	8.610
5	0.727	1.476	2.015	2.571	3.365	4.032	6.859
6	0.718	1.440	1.943	2.447	3.143	3.707	5.959
7	0.711	1.415	1.895	2.365	2.998	3.499	5.405
8	0.706	1.397	1.860	2.306	2.896	3.355	5.041
9	0.703	1.383	1.833	2.262	2.821	3.250	4.761
10	0.700	1.372	1.812	2.228	2.764	3.169	4.587
11	0.697	1.363	1.796	2.201	2.718	3.106	4.437
12	0.695	1.356	1.782	2.179	2.681	3.055	4.318
13	0.694	1.350	1.771	2.160	2.650	3.012	4.221
14	0.692	1.345	1.761	2.145	2.624	2.977	4.140
15	0.691	1.341	1.753	2.131	2.602	2.947	4.073
16	0.690	1.337	1.746	2.120	2.583	2.921	4.015
17	0.689	1.333	1.740	2.110	2.567	2.898	3.965
18	0.688	1.330	1.734	2.101	2.552	2.878	3.922
19	0.688	1.328	1.729	2.093	2.539	2.861	3.883
20	0.687	1.325	1.725	2.086	2.528	2.845	3.850
21	0.686	1.323	1.721	2.080	2.518	2.831	3.819
22	0.686	1.321	1.717	2.074	2.508	2.819	3.792
23	0.685	1.319	1.714	2.069	2.500	2.807	3.767
24	0.685	1.318	1.711	2.064	2.492	2.797	3.745
25	0.684	1.316	1.708	2.060	2.485	2.787	3.725
26	0.684	1.315	1.706	2.056	2.479	2.779	3.707
27	0.684	1.314	1.703	2.052	2.473	2.771	3.690
28	0.683	1.313	1.701	2.048	2.467	2.763	3.674
29	0.683	1.311	1.699	2.045	2.462	2.756	3.659
30	0.683	1.310	1.697	2.042	2.457	2.750	3.646
40	0.681	1.303	1.684	2.021	2.423	2.704	3.551
60	0.679	1.296	1.671	2.000	2.390	2.660	3.460
120	0.677	1.289	1.658	1.980	2.358	2.617	3.373
∞	0.674	1.282	1.645	1.960	2.326	2.576	3.291

* Taken from Table III of R. A. Fisher and F. Yates: "Statistical Tables for Biological, Agricultural, and Medical Research," published by Oliver & Boyd Ltd., Edinburgh, by permission of the authors and publishers.

